

IN THE CLAIMS**COMPLETE LISTING OF ALL CLAIMS, WITH MARKINGS AND STATUS IDENTIFIERS**

In the revised claim set appearing below, currently amended claims have deletions shown by ~~strike through~~ or brackets [[]], and additions shown by underlining. This listing of claims will replace all prior versions and listings of the claims in the application.

Listing of Claims:

1. (canceled)
2. (currently amended) The method ~~A peptide~~ according to claim [[1]] 6 or 9, wherein said peptide of formula (I) is a peptide wherein
 - A¹ is L-Phe, D-Phe, L-Cpa or D-Cpa;
 - A³ is L-Tyr, L-Trp or L-3-Pal;
 - A⁴ is D-Trp;
 - A⁶ is β -Ala or Gaba;
 - A⁷ is L-Cys;
 - A⁸ is L-Thr, L-Trp, L-Leu or L-Nal; and
 - R² and R³ are each H;
 - or a pharmaceutically acceptable salt thereof.
3. (currently amended) The method ~~A peptide~~ according to claim [[2]] 6 or 9, wherein said peptide is of the formula
 - Cpa-cyclo(D-Cys-3-Pal-D-Trp-Lys-Gaba-Cys)-Nal-NH₂;
 - Cpa-cyclo(D-Cys-3-Pal-D-Trp-Lys- β -Ala-Cys)-Nal-NH₂;
 - Phe-cyclo(D-Cys-3-Pal-D-Trp-Lys-Gaba-Cys)-Nal-NH₂;
 - Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Gaba-Cys)-Nal-NH₂;
 - Phe-cyclo(D-Cys-Trp-D-Trp-Lys-Gaba-Cys)-Nal-NH₂;
 - Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Gaba-Cys)-Trp-NH₂;
 - D-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Gaba-Cys)-Nal-NH₂;


D-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Gaba-Cys)-Leu-NH₂; or
 Phe-cyclo-(D-Cys-Tyr-D-Trp-Lys-Gaba-Cys)-Thr-NH₂;
 or a pharmaceutically acceptable salt thereof.

4. (currently amended) The method A-peptide according to claim [[3]] 6 or 9, wherein said peptide is of the formula

Cpa-cyclo(D-Cys-3-Pal-D-Trp-Lys-Gaba-Cys)-Nal-NH₂; or
 Cpa-cyclo(D-Cys-3-Pal-D-Trp-Lys-β-Ala-Cys)-Nal-NH₂;
 or a pharmaceutically acceptable salt thereof.


5. (currently amended) [[A]] The method according to claim 6 or 9, wherein said peptide or pharmaceutically acceptable salt thereof is in the form of a pharmaceutical composition useful for eliciting a somatostatin agonist response in a human or other animal which comprises an effective amount of a peptide of formula (I) ~~according to claim 1~~ or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

6. (previously presented) A method of eliciting a somatostatin agonist response in a human or other animal in need thereof, which comprises administering an effective amount of a peptide of formula (I)



(I)

or a pharmaceutically acceptable salt thereof,
 wherein



A¹ and A³ are each independently the D- or L-isomer of an amino acid selected from the group consisting of Phe, Tyr, Tyr(I), Trp, 3-Pal, 4-Pal, Cpa and Nal;

A⁴ is L-Trp, D-Trp, L-β-methyl-Trp or D-β-methyl-Trp;

A⁶ is -NH-(CHR¹)_n-CO-, where n is 2, 3, or 4;

A⁷ is L- or D-Cys;


A⁸ is the D- or L-isomer of an amino acid selected from the group consisting of Phe, Tyr, Tyr(I), Trp, Nal, Cpa, Val, Leu, Ile, Ser and Thr;

Y is NR^2R^3 where R^2 and R^3 are each independently H or $(\text{C}_1\text{-C}_5)\text{alkyl}$;

R^1 is selected from the group consisting H, $(\text{C}_1\text{-C}_4)\text{alkyl}$ and $-\text{CH}_2\text{-aryl}$; wherein said aryl is an optionally substituted moiety selected from the group consisting of phenyl, 1-naphthyl, and 2-naphthyl, wherein said optionally substituted moiety is optionally substituted with one or more substituents each independently selected from the group consisting of $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{2-6})\text{alkenyl}$, $(\text{C}_{2-6})\text{alkynyl}$, aryl, aryl $(\text{C}_{1-6})\text{alkyl}$, $(\text{C}_{1-6})\text{alkoxy}$, $-\text{N}(\text{R}^4\text{R}^5)$, $-\text{COOH}$, $-\text{CON}(\text{R}^4\text{R}^5)$, halo, $-\text{OH}$, $-\text{CN}$, and $-\text{NO}_2$;

R^4 and R^5 each is, independently for each occurrence, H or $(\text{C}_{1-3})\text{alkyl}$;
where the Cys of A^2 is bonded to the Cys of A^7 by a di-sulfide bond formed from the thiol groups of each Cys,
to the human or other animal.


7. (canceled)
8. (canceled)
9. (currently amended) A method of inhibiting the secretion of growth hormone, insulin, glucagon or pancreatic exocrine secretion in a human or other animal in need thereof, which comprises administering a peptide of formula (I)



(I)

or a pharmaceutically acceptable salt thereof,

wherein



A^1 and A^3 are each independently the D- or L-isomer of an amino acid selected from the group consisting of Phe, Tyr, Tyr(I), Trp, 3-Pal, 4-Pal, Cpa and Nal;

A^4 is L-Trp, D-Trp, L- β -methyl-Trp or D- β -methyl-Trp;

A^6 is $-\text{NH}-(\text{CHR}^1)_n\text{-CO-}$, where n is 2, 3, or 4;

A^7 is L- or D-Cys;

A⁸ is the D- or L-isomer of an amino acid selected from the group consisting of Phe, Tyr, Tyr(I), Trp, Nal, Cpa, Val, Leu, Ile, Ser and Thr;

Y is NR²R³ where R² and R³ are each independently H or (C₁-C₅)alkyl;

R¹ is selected from the group consisting H, (C₁-C₄)alkyl and -CH₂-aryl; wherein said aryl is an optionally substituted moiety selected from the group consisting of phenyl, 1-naphthyl, and 2-naphthyl, wherein said optionally substituted moiety is optionally substituted with one or more substituents each independently selected from the group consisting of (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, aryl, aryl(C₁₋₆)alkyl, (C₁₋₆)alkoxy, -N(R⁴R⁵), -COOH, -CON(R⁴R⁵), halo, -OH, -CN, and -NO₂;

R⁴ and R⁵ each is, independently for each occurrence, H or (C₁₋₃)alkyl;
where the Cys of A² is bonded to the Cys of A⁷ by a di-sulfide bond formed from the thiol groups of each Cys,
to said human or other animal.

10. (canceled)

11. (canceled)